

Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at http://about.jstor.org/participate-jstor/individuals/early-journal-content.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

liquity, gives the diminution for an interval of nearly sixty years, with almost sufficient accuracy to state with some confidence the mass of Venus; but to obtain this point with certainty, the present obliquity, deduced from a mean of the observations of different astronomers, should be used. Upon this subject the author alludes to the opinion of astronomers, that observations of the winter solstice have given a less obliquity than those of the summer solstice,—an opinion sustained by the observations of Maskelyne, Arago, and Pond, but questioned by Bessel and Bradley. Dr. Brinkley refers this difference to some unknown modification of refraction; he has observed that at the winter solstice the irregularity of refraction for the sun is greater than for the stars at the same zenith-distance. He points out the necessity of paying attention to the observations at the winter solstice, and gives a table, exhibiting the mean obliquity reduced to January 1813.

Dr. Brinkley next alludes to the maximum of the aberration of light, which appears from his observations of last year to be 20".80.

On some New Methods of investigating the Sums of several Classes of Infinite Series. By Charles Babbage, Esq. A.M. F.R.S. Read April 1, 1819. [Phil. Trans. 1819, p. 249.]

The object of this paper is to explain two methods of finding the sums of a variety of infinite series. One of these the author discovered several years ago; but finding that some of the results to which it led were erroneous, he then declined publishing it. In inquiring into the causes of these errors, he was led to the second method, which employs the process of integration relative to finite differences. The cause of the fallacies in the former method was afterwards discovered, and in this paper a criterion is proposed for judging of the truth of the results, and a mode of correcting them where found to be erroneous. The sums of a variety of series are found by these methods; and the author concludes by observing, that he has since been informed by M. Poisson, that that gentleman had arrived at some nearly similar results in investigating a problem in physical astronomy, and also that some investigations of a similar nature were found amongst the papers of Lagrange, but that neither of these mathematicians had explained the cause of the errors, or given a method of correcting them.

On the Optical and Physical Properties of Tabasheer. By David Brewster, LL.D. F.R.S. Lond. and Edin. In a Letter to the Right Hon. Sir Joseph Banks, Bart. G.C.B. P.R.S. &c. Read May 6, 1819. [Phil. Trans. 1819, p. 283.]

Tabasheer is a substance found in the cavities of the bamboo, existing originally in the state of a transparent fluid, but gradually indurating into a solid of different degrees of hardness: it consists of 70 silica, +30 potash and lime. One variety has a milky trans-